



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

volve an infinite acquired velocity, which is absurd.' In this way Peirce justifies Galileo's conclusion.

The error in Peirce's reasoning seems to me perfectly apparent. When $s=0$ he takes C to be an infinitesimal, while really C is an absolute zero. When s and t are both zero, e^{at} is not zero, hence e^{at} , multiplied by an infinitesimal, cannot be equal to absolute zero. An infinitesimal is a variable whose limit is zero, but the variable never reaches its limit. If we consider an infinitesimal an extremely small quantity, we must still remember that it is a quantity. Now, if C is absolute zero, then s can never be different from zero, no matter how large e^{at} may be.

It is easy to illustrate my conclusion by physical examples. A particle is placed in a smooth tube which revolves horizontally about an axis through its center. With what velocity will the particle move? The only force impelling the particle along the tube is the centrifugal force due to rotation. Hence we have $\frac{d^2s}{dt^2}=w^2s$ and $\frac{ds}{dt}=ws$, where w is the uniform angular velocity. Here the velocity is proportional to the distance from the axis. Suppose now that the particle lies initially at rest in the axis. Will it begin to move? There is no reason why it should move one way any more than the other.

The two assumptions in our problem are contraries. The first excludes the possibility of motion; the second declares that motion exists. From assumptions that are contraries no conclusion can be drawn.

DIOPHANTINE ANALYSIS.

80. Proposed by M. A. GRUBER, A. M., War Department, Washington, D. C.

Find three square numbers whose reciprocals form an arithmetical progression.

I. Solution by the PROPOSER.

When three numbers are in arithmetical progression, the products of these numbers, taken two at a time, will give three numbers whose *reciprocals* are in arithmetical progression.

Let $x-a$, x , and $x+a$ =three numbers in arithmetical progression. Then $x(x-a)$, $(x-a)(x+a)$, and $x(x+a)$ =the three numbers whose reciprocals are in arithmetical progression.

$$\text{For, } \frac{1}{x(x-a)} - \frac{1}{(x-a)(x+a)} = \frac{1}{(x-a)(x+a)} - \frac{1}{x(x+a)} = \frac{a}{x(x-a)(x+a)}.$$

The general values for three *squares* in arithmetical progression are $(p^2 - q^2)^2$, $2pq^2$, $(p^2 + q^2)^2$, and $(p^2 - q^2 + 2pq)^2$, where $x = (p^2 + q^2)^2$, and a the common difference $= 4pq(p^2 - q^2)$.

Put $p=2$ and $q=1$; then 1^2 , 5^2 , and 7^2 are three squares in arithmetical progression. Whence the three squares whose reciprocals form an arithmetical

